

The Entomologist's Record

To encourage the publication of concise and useful new distribution records, corrections of previously published erroneous records, misidentifications, short field notes, and current news items about Entomologists, amateur and professional, Entomology Departments and Museums, prompt publication is offered in this Department.

NEW RECORDS FOR TRICHOPTERA IN NORTH DAKOTA¹

S.C. Harris, R.B. Carlson²

ABSTRACT: The known ranges of *Cheumatopsyche pasella* Ross and *Hydropsyche piatrix* Ross are extended westward to include North Dakota.

DESCRIPTORS: Hydropsychidae, State records, North Dakota.

Cheumatopsyche pasella Ross is primarily northeastern in distribution, extending as far south as Georgia and as far west as Wisconsin (Miller, 1965). *C. pasella* was collected in Richland and Ransom Counties, North Dakota during an investigation of small sandhill streams. A series of 54 adults, 12 males and 42 females, were collected in light traps in June and July, 1974.

Ross (1944) gives the distribution of *Hydropsyche piatrix* Ross as Missouri and Arkansas. Two females of this species were collected in light traps placed on the banks of a small stream in Richland County, North Dakota. The specimens were collected on June 5 and 9, 1974.

The assistance of Dr. Andrew Nimmo, Department of Entomology, University of Alberta, in identifying the specimens, is gratefully acknowledged.

LITERATURE CITED

- Miller, R.B. 1965. The taxonomy and distribution of the genus *Cheumatopsyche* in the northeastern United States (Trichoptera: Hydropsychidae). Ph.D. Thesis, Pennsylvania State University.
- Ross, H.H. 1944. The caddisflies, or Trichoptera, of Illinois. Ill. Nat. Hist. Surv. Bull. 23: 1-326.

¹Accepted for publication: March 12, 1977

²Department of Entomology, North Dakota State University, Fargo, North Dakota 58102